

摘 要

北海道の秋播小麦にとって、耐凍性の問題は古くからの重要な育種目標であるが、特性が複雑な上、各種の雪腐病抵抗性が関与するため、積極的な育種をはばんできた。本報告は耐凍性の要因中、耐凍性と雪腐病抵抗性および春化要求度（秋播性程度）を研究の対象としたが、とくに耐凍性、雪腐大粒菌核病および雪腐小粒菌核病抵抗性の3要因を育種的にどこまで高めうるかを確かめようとして行われた。えられた結果を要約するとつぎのとおりである。

1. 耐凍性の品種間差異

1) 雪腐大粒菌核病の検定には、小麦を20cmの高畦で栽植し、11月中旬に子のう孢子散布を行い、さらに低濃度の薬剤防除と無防除の区を設けることによって、被害の激甚な年にも軽微な年にも確実な検定が可能となった。

2) 雪腐小粒菌核病および紅色雪腐病の検定にはBruehlらの大量培養法に準じて病原菌を培養し、苗箱に栽植した小麦に接種した。品種間差異が明瞭になるのは、接種後2℃の低温恒温室に置いて、黒色小粒菌核病で50~60日、褐色小粒菌核病で90日、紅色雪腐病では60日の処理期間を要した。この方法による検定結果は多雪地帯のは場検定との間に有意な相関を示した($r=0.555^{**} \sim 0.633^{***}$)。

3) 耐凍性の検定には地表露出法と冠部凍結法およびE.C.法を検討し、いずれも精度は高く、前2者間の相関は $r = 0.733^{***}$ で、後2者間の相関は $r = 0.768^{***}$ であった。E.C.法でみると、訓子府において11月中旬まで低温順化された小麦は、 -11.5°C から凍結被害が現れ、 -18°C 以下になると強品種でも被害が増大する。 -15°C が凍結処理温度として最適と見られた。

4) それぞれの検定結果、次の有用母材が見いだされたが、いずれの品種も農学実用形質は貧弱である。

S. borealis 抵抗性：「Iohardi」「北海48号」。

T. ishikariensis 抵抗性：「C. I. 14106」、
「P. I. 172582」、「P. I. 173438」。

T. incarnata 抵抗性：「C. I. 14106」、
「P. I. 173438」、「農林62号」、「北系 628」。

F. nivale 抵抗性：「C. I. 14106」「P. I. 172582」、
「P. I. 173438」。

耐凍性：「Lutescens 0329」、「Valujevskaja」。

5) 耐凍性品種は *S. borealis* におおむね抵抗性を示したが ($r=0.443^{*} \sim 0.550^{**}$)、「Lutescens 0329」、「Valujevskaja」はきわめて耐凍性でありながら、*S. borealis* にはあまり強くないし、*S. borealis* に強い「北海48号」の耐凍性はむしろ弱いなどの特異な反応を示す品種が認められた。

6) *Typhula* spp. および *F. nivale* 抵抗性と耐凍性あるいは *S. borealis* 抵抗性との間には、弱い負の相関が認められ、*T. ishikariensis*、*T. incarnata* および *F. nivale* 抵抗性3者間には高い正の相関がみられた ($r = 0.555^{**} \sim 0.799^{***}$)。

7) どの要因にもすべて抵抗性という品種は認められず、上記の関係から25品種系統を4群に分類することができた (Fig. 3)。

A群 (非耐凍型)：「Ibis」、「Gaines」、「タクネコムギ」。

B群 (耐凍型)：「Lutescens 0329」ほか8品種。

C群 (中間型)：「ホロシリコムギ」ほか道内の9品種・系統。

D群 (耐雪型、*Typhula* spp. 抵抗性)：「C. I. 14106」、「P. I. 172582」、「P. I. 173438」、「農林62号」、「北系 628」。

2. 耐凍性の遺伝

1) *S. borealis*、*T. ishikariensis*、*T. incarnata* および凍害による被害度の平均値と分散は、いずれも組合せ間、親品種間に有意差が認められ、各 F_2 集団の平均値と分散はいずれも中間親の値に左右された。

- 2) ダイアレル分析の前提となる6条件のうち、抵抗性の遺伝行動が非対立遺伝子間の相互作用がないことを検定し、エピステーシス効果のない単純な相加一優性モデルが想定された。
- 3) 分析の結果、各要因とも相加的遺伝子効果が高かった。*S. borealis* 抵抗性は部分優性、*T. ishikariensis* 抵抗性と耐凍性は超優性であった。*S. borealis* と *T. ishikariensis* 抵抗性は劣性遺伝子が支配的なものに対し、耐凍性では一定の傾向が認められなかった。
- 4) 狭義の遺伝力は *S. borealis* 抵抗性：90、*T. incarnata* 抵抗性：79、耐凍性：82%と高く推定された。また、 F_2 、 F_3 集団の親子相関による遺伝力は *S. borealis* 抵抗性：84、*T. ishikariensis* 抵抗性：48%と推定された。
- 5) $K = (\text{最大の親の値} - \text{最小の親の値})^2 / 4D$ による有効因子数は、*S. borealis* 抵抗性：1.88、*T. ishikariensis* 抵抗性：2.32、耐凍性：2.03と推定され、関与する遺伝子数は2～3個以上と考えられた。
- 6) 遺伝相関 (r_G) は凍害と *S. borealis* : 0.520、凍害と *T. ishikariensis* : -0.819、*S. borealis* と *T. ishikariensis* : -0.818と推定され、耐凍性に関与する遺伝子の中には *S. borealis* の抵抗性に関与する遺伝子が含まれていることが示唆されたのに対し、これらと *T. ishikariensis* 抵抗性に関与する対立遺伝子は互いに相異なるように考えられた。
- 7) *S. borealis* と *T. ishikariensis* に対して、無被害個体の出現率を F_2 集団で比較すると、全体として $r = -0.592^{**}$ と負の関係が有意であるが、両病害に抵抗性個体を多く含み、耐凍性も強い有望な4組合せが認められ、このうち3組合せは道内品種・系統間の交雑であった。道内品種・系統がもつ耐冬性の遺伝的背景の重要性が示唆された。

3. 耐冬性と体内成分の関係

1) 地上部の乾物重、全炭水化物、糖類、粗デンプンおよび蛋白態窒素は秋から初冬まで急速に増加し、根雪直後最高に達する。可溶性窒素、リン酸、カリ、ケイ酸および灰分は秋季低温とともに

減少する。

- 2) 越冬前に測定した16形質では地域、時期、品種および越冬型の間に有意な差異が認められた。
- 3) 訓子府は岩見沢より生育量は小さいが、水分が少なく、炭水化物や糖の蓄積が多く、その消耗も少ない。
- 4) *Typhula* spp.と *F. nivale*に抵抗性の品種は越冬前に多量の炭水化物を蓄積し、越冬後にもなお多量の炭水化物を維持した。また越冬後の可溶性窒素は著しく少なかった。
- 5) 越冬前と越冬後の体内成分の相関は還元糖と粗デンプンを除くといずれも有意に高かった。全炭水化物の年次間相関は訓子府で越冬前 0.751^{***} 、越冬後 0.767^{***} と高く、年次が異なっても品種間差異はほぼ平行的であった。
- 6) 凍害と浸透価、*S. borealis* とリン酸および脂質の間にきわめて高い相関がみられ、*T. ishikariensis*、*T. incarnata* および *F. nivale* に共通して最も高い相関を示したのは蛋白態窒素と全炭水化物で、越冬前・後ともに蛋白態窒素が多いと被害が増大する。
- 7) 植物体の生育量および越冬前後の体内成分の22形質を用いて主成分分析を行い、品種の生態的特性を4群に分類することができた (Fig.7)。このグループは第1章の越冬型の分類に非常によく適合した。
- 8) *Typhula* spp.および *F. nivale*に抵抗性の「C. I. 14106」、「P. I. 172582」および「P. I. 173438」は、他の越冬型の品種より炭水化物や糖の蓄積がきわめて多く、逆に蛋白態窒素、リン酸、脂質が少なく、浸透価が著しく低かった。このような事実は、これら3品種が凍害に弱く *S. borealis* に罹病性であることと関係し、特異な遺伝的特性に基づくものと思われた。

4. 耐冬性と春化要求度の関係

1) 緑体春化法により春化所要日数を推定すると、北海道の品種系統は40日の「タクネコムギ」から70日の「北海48号」まで分布し、さらに90日あるいは100日を要すると思われる外国品種が存在した。春化要求度に基づき、供試品種を4つのグル

ープに群別した。

2) 第1グループの「タクネコムギ」は春化要求度Ⅵに相当するが、Ⅶに相当する品種群は広い変異を示したので、「ムカコムギ」、「ホクエイ」で代表される第2グループ(春化所要日数50~60日)と「赤錆不知1号」、「農林8号」(同65日)の第3グループに分け、「北海48号」もこれに含めた。第4グループは60日低温処理では完全に出穂しないもので、耐凍性のソビエト品種「Moscow 1」、「Valujevskaja」、「Lutescens0329」と、*Typhula* spp. 抵抗性の「P. I. 172582」、「P. I. 173438」、「C. I. 14106」の6品種であった。

3) これら品種の長期間の春化処理には、8℃の緑体春化より晩秋戸外の低温短日処理が効果的であった。また第4グループ品種は、従来の播種Ⅶないし春化要求度Ⅶをはるかに越えるものであり、Ⅷ以上に相当するものと考えられた。

4) 春化要求度と耐凍性との関係では、密接な品種群と、小数ながら無関係の品種群が存在した。*T. ishkariensis* および *S. borealis* 抵抗性とは有意な関係は認められなかった。

5) 「P. I. 173438」および「Valujevskaja」と「チホクコムギ」との交雑後代F₄系統では、出穂まで日数の分布幅が45日におよび、緑体春化区戸外区を用いて5つに群別が可能であった。このことから両親品種の春化要求度の差異には複数の遺伝子が関与することが暗示された。

6) 「Valujevskaja」に近い耐凍性系統、あるいは「P. I. 173438」並の *T. ishkariensis* 抵抗性が第1、3、4の各グループから同程度の頻度で見いだされたこと、また春化要求度との間に有意な相関が認められなかったことから、春化要求度の高い段階ではこれらの特性はそれぞれ独立と考えられた。

5. 耐凍性の選抜実験

1) *T. ishkariensis* B型のほ場での接種検定は、やや区間変動が大きく精度に難点があるが、同病による単一発生がえられ、選抜法として有効と考えられた。

2) 「P. I. 173438」／「ホロシリコムギ」／「北見42号」のF₆ 91群 364系統の接種検定の結果、「P. I. 173438」並の抵抗性を示すものが13系統群認められた。いずれも群間の変異係数も小さく、十分な *T. ishkariensis* 抵抗性を持つと判断された。これらのうち、稈長、強稈性が「チホクコムギ」に近く、千粒重が「チホクコムギ」より重く、収穫指数が「ホロシリコムギ」以上の系統「WH 7, 8, 9, 10」は農学的特性が優れ有望とみられた。

3) 耐凍性と *T. ishkariensis* 抵抗性の同時選抜を実施した「ホロシリコムギ」／「P. I. 173438」／「Valujevskaja」／「ホロシリコムギ」のF₈ 980系統からは「Valujevskaja」並の耐凍性を持つものが91系統えられ、「P. I. 173438」並の *T. ishkariensis* 抵抗性系統は76えられた。さらに両者に対して抵抗性の系統は6系統で全系統の0.6%であった。農学的特性は「ホロシリコムギ」に比較すると劣るが、「P. I. 173438」と「Valujevskaja」の改良型として有望とみられた。

4) *S. borealis*, *Typhula* spp. 抵抗性および耐凍性の3要因を同時に満足する系統は得られなかった。「Moscow 1」／「ホロシリ」のF₇で *S. borealis* に抵抗性の超越分離がみられ、そのうち「WH 11, 12」の2系統は「Moscow 1」並の耐凍性を示した。ただし農学的特性はなお若干の改良を要する。

5) 耐凍性と *Typhula* spp. 抵抗性およびその両者については、以上の系統を用いた育種計画で実用品種の育成は可能と判断されたが、*S. borealis* 抵抗性育種については遺伝子源が乏しく、将来の課題として残された。

引用文献

- 1) 阿部二郎 “イネ科牧草の耐寒性に関する品種間差異” 日草誌. 25, 279-284 (1980)
- 2) 阿部直隆 “麦類雪腐病に対するチオファネートメチル剤の効果について” 北農40(5), 30-41(1973)
- 3) Adair, C. R. and J. W. Jones “Effect of environment on the characteristics of plants surviving in bulk hybrid populations of rice” J. Amer. Soc. Agron. 38, 708-716(1946).
- 4) Allan, R. E. and J. A. Pritchett “Inheritance and association of subcrown-internode length with coleoptile and culm length in a winter wheat cross” Crop Sci. 13, 639-641 (1973).
- 5) Amirshahi, M. C. and F. L. Patterson “Development of a standard artificial freezing technique for evaluating cold resistance of oats” Agron. J. 48, 181-184 (1956).
- 7) Anderson A. and T. A. Kiesselbach “Studies on the technique of controlled hardiness tests with winter wheat” J. Amer. Soc. Agron. 26, 44-50 (1934).
- 8) Andrews, J. E. “Controlled low temperature tests of sprouted seeds as a measure of cold resistance of winter wheat varieties” Canad. J. Plant Sci. 38, 1-7 (1958).
- 9) Årsvoll, K. “Effests of hardening, plant age, and development in *Phleum pratense* and *Festuca pratensis* on resistance to snow mold fungi” Meld. Norg. LandbrHøgsk. 56(28), 1-14 (1977).
- 10) Årsvoll, K. “Effects of nitrogen, phosphorus, and potassium on resistance to snow mold fungi and on freezinfg tolerance in *Phleum platense*” Meld. Norg. LandbrHøgsk. 56 (29), 1-14 (1977)
- 11) Ausemus, E. R. , F. H. McNeal and J. W. Schmidt “Genetics and inheritance. In Wheat and wheat improvemint” Amer. Coc. Agron. Inc. , Madison, Wis. Press, Agronomy 13 (1967)
- 12) Bruehl, G. W. , R. Sprague, W. R. Fischer, M. Nagamitsu, W. L. Nelson, and O. . A. Vogel, “Snow mold of winter wheat in Washington” Wah. Agr. Exp. Stn. Bull. 677, 1-21 (1967).
- 13) Bruehl, G. W. “Effect of plant size on resistance to snow mold of winter wheat” Plant Dis. Rep. 51, 815-819 (1967).
- 14) Bruehl, G. W. , M. Nagamitsu, W. L. Nelson, C. J. Peterson, Jr. and G. L. Rubenthaler “Registration SPRAGUE wheat” Crop Sci. 18, 695-696. (1978).
- 15) Bruehl, G. W. “Development wheats resistant to snow mold in Washington State” Plant Disease Rep. 66, 1090-11095 (1982).
- 16) Bruehl, G. W. “Correlation of resistance to *Typhula idahoensis*, *T. incarnata*, and *Fusarium nivale* in certain varieties of winter wheat” Phytopathology57 (3), 308-310 (1967)
- 17) Bruehl, G. W. and B. Cunfer “Physiologic and environmental factors that affect the severity of snow mold of wheat” Phytopath. 661 (7), 792-799 (1971)
- 18) Bruehl, G. W. , R. Kiyomoto, C. Peterson and M. Nagamitsu “Testing winter wheats for snow mold resistance in Washington” Plant Dis. Rep. 59, 566-570 (1875).
- 19) Cahalan, C. and C. N. Law “The genetical control of cold resistance and vernalization requirement in wheat” Heredity 42, 125-132 (1979).

- 20) Chen, P. and L. V. Gusta "The role of water in cold hardiness of winter cereals. In Plant cold hardiness" Li, P. H. and A. Sakai Ed. Academic Press, 165-174 (1978).
- 21) Christiansen, M. N. and C. F. Lewis "Breeding plants for less favorable environments" John Wiley & Sons. Press, Inc. (1982).
- 22) Coffman, F. A. "Winter oats extended northward by breeding and testing" *Agron. J.* 49, 187 (1957).
- 23) Compton, L. E. , Caldwell, R. M. , J. F. Sokafer and F. L. Patterson "Norline winter oats" *Purdue Univ. Agric. Res. Stn. Res. Bull. No. 799*, 6 p. (1965).
- 24) Cormack, M. W. and J. B. Lebeau "Snow mold infection of alfalfa, grasses and several fungi under artificial conditions" *Can. Bot.* 37, 685-693 (1959)
- 25) Dexter, S. T., W. E. Tottingham, and L. F. Graber "Investigation of the hardiness of plants by measurement of electrical conductivity" *Plant Physiol.* 7, 63-78 (1932).
- 26) Dexter, S. T. "Effect of several environmental factors on the hardening of plants" *Plant Physiol.* 8, 123-139 (1933).
- 27) Dexter, S. T. "The evaluation of crop plants for winter hardiness" *Adv. Agron.* (, 203-239. (1956).
- 28) Dorofeev, V. F. "Evaluation of material for frost and drought resistance in wheat breeding" *IBP. No. " Camb. Univ. Press*, 211-222 (1975).
- 29) Dvorak, G. and D. B. Fowler "Cold hardiness potential of Triticale and Tetraploid rye" *Crop Sci.* 18, 477-478 (1978).
- 30) Erickson, J. R. , L. D. Sibbitt and J. D. Miller "Registration of Roughrider wheat" *Crop Sci.* 17, 980 (1977).
- 31) Eunus, A. M. , Johnson, L. P. and R. Aksil "Inheritance of winter hardiness in an eighteen-parent diallel cross in barley" *Can. J. Genet. Cytol.* 4, 356-377 (1962).
- 32) Finkner, V. C. "Effect of natural selection on winter survival of winter oat bulk hybrid populations" *Crop Sci.* 4, 465-466 (1964).
- 33) Finkner, V. C. , D. L. Davis, C. R. Tutt and J. T. Green "Registration of Walkin winter oat" *Crop Sci.* 11, 133 (1971).
- 34) Flood, R. G. and G. M. Halloran "Basic development rate in spring wheat" *Agron. J.* 76, 260-264 (1984).
- 35) Flood, R. G. and G. M. Halloran "The nature and duration of gene action for vernalization response in wheat" *Ann. Bot.* 53, 363-368 (1984).
- 36) Florell, V. H. "Bulked population method of handling cereal hybrids" *J. Amer. Soc. Agron.* 21, 718-724 (1929).
- 37) Fowler, D. B. , J. Dvorak, and L. V. Gusta "Comparative cold hardiness of several *Triticum* species and *Secale cereale* L. " *Crop Sci.* 17, 941-943 (1977).
- 38) Fowler, D. B. and L. V. Gusta "Selection for winterhardiness. I. Identification of genotypic variability" *Crop Sci.* 19, 769-772 (1979).
- 39) Fowler, D. B. "Selection for winterhardiness. II. Variation within field trials" *Crop Sci.* 19, 773-775 (1979).

- 40) Fowler, D. B. , L. V. Gusta and N. J. Tyler “Selection for winterhardiness in wheats. III. Screening methods” *Crop Sci.* 21, 896-901 (1981).
- 41) Fowler D. B. and R. J. Carles “Growth, development, and cold tolerance of fall acclimated cereal grains” *Crop Sci.* 19, 915-922 (1979).
- 42) 後藤寛治 “育種上における自然淘汰の意義” *遺伝*7 (5,6), (1953).
- 43) 後藤寛治 “育種へのいざない, 研究遍歴の道程” *北農研究シリーズ*V. 206p (1973).
- 44) 後藤虎男 “コムギの出穂性に関する生態学的ならびに遺伝学的研究” *東北農試研報.* 59. 1-69 (1979).
- 45) Gotoh, T. “Variational variation and inheritance mode of vernalization and inheritance mode of vernalization requiremint in common wheat” *Proc. 6th. Int. Wheat Gen. Sympo.* , 475-478 (1983).
- 46) Grafius, J. E. “Winter hardiness in barley. In *Breeding for resistance*” *Michigan Agric. Exp. Stn. Res. Rep.* 247, 16 (1974).
- 47) Grafius, J E. “Breeding for winter hardiness. In *Analysis and improvement of plant cold hardiness.*” *CRC. Press Inc.* 161-174 (1981).
- 48) Granhall, I. “Genetic and physiological studies in interspecific wheat crosses” *Hereditas* 29, 269-373 (1975).
- 49) Gullord, M. , C. R. Olien and E. M. Everson “Evaluation of freezing hardiness in winter wheat” *Crop Sci.* 15, 153-157 (1975).
- 50) 北海道経済部. 北海道農業試験場共編 “麦類雪腐病の防除法について” *農事指導資料.* (1949).
- 51) 北海道農業試験場作物部作物第2研究室 “麦類の耐冬性に関する研究” *昭和33年度試験成績書.* (1958).
- 52) 北海道立農業試験場北見支場 “耐冬性に関する研究” *昭和30年度秋播麦類試験成績書*(1956).
- 53) 北海道立農業試験場北見支場 “耐冬性に関する研究” *昭和31年度秋播麦類試験成績書*(1956).
- 54) Harlan, H. V. , M. L. Martini and H. Stevens “A study of method in barley breeding” *USDA Tech. Bull. Mo.* 720, 26 pp. (1940)
- 55) Hayes, H. K. and O. S. Aamodt “Inheritance of winter hardiness and growth habit in crosses of Marquis with Minhardi and Minturki wheats” *J. Agric. Res.* 35, 223-236 (1927)
- 56) Hayman, B. I. “The theory and analysis of diallel crosses” *Genetics* 39, 789-809 (1954)
- 57) Hayman, B. I. “The theory and analysis of diallel crosses” *Biometrics* 10, 235-244 (1954)
- 58) Hill, D. D. and S. C. Salmon “The resistance of certian varieties of winter wheat to artificially produced low temperature” *J. Agric. Res.* 35, 933-937 (1927)
- 59) Hirai Tokuzo “Studies on the nature of disease resistance in plants. A contribution to the knowledge concerning the mechanisms of infection of plants by the weak pathogenic fungi under snow cover and the nature of disease resistance against fungi or viruses on the basis of histo and cytochemical observations” *Forsch. Gebeit. Pflanzenkr. (Kyoto)* 5, 139-157 (1956)
- 60) 広川文彦 “PCPによる冬損防止と雑草防除” *PCP協議会資料.* 7, (1963)
- 61) 広川文彦 “麦類の耐冬性検定法に就いて” *日本育種. 作物. 遺伝学会北海道談話会講演要旨.* (1960)
- 62) Hoogendoorn, J. “The basis of variation in data of ear emergence under field conditions among the progeny of a cross between two winter wheat varieties” *J. Agric. Sci.* 104, 493-500 (1985)
- 63) Hoogendoorn, J. “Basic development rate in spring wheat” *Agron. J.* 76, 260-264 (1984)

- 64) Jamaleinen, E. A. "Resistance in winter cereals and grasses to low-temperature parasitic fungi" *Ann. Rev. Phytopath.* 12, 281-302 (1974)
- 65) Jinks, J. L. "The F₂ and backcross generations from a set of diallel crosses" *Heredity* 10, 1-30 (1956)
- 66) Jones, R. M. "Analysis of variance of the half diallel table" *Heredity* 20, 177-121 (1965).
- 67) 柿崎洋一, 鈴木真三郎 "小麦における出穂の生理に関する研究" *農事試験報*, 3, 41-91 (1937)
- 68) 木村幹男, 平井篤造 "ムギ類雪腐病に関する研究. 第6報. ムギ類品種の抵抗性検定方法" *東北農業*. 5 (5. 6) 259-261 (1952)
- 69) Kiyomoto, R. K. and G. W. Bruehl "Carbohydrate accumulation and depletion by winter cereals differing in resistance to *Typhula idahoensis*" *Phytopathology* 67, 206-210 (1977)
- 70) Kraimi, Y. Y. and C. O. Qualset "Genetics of heading time in wheat. II. The inheritance of vernalization response" *Genetics* 76, 119-133 (1974)
- 71) 国井輝男 "麦類雪腐病に関する既往の研究について" *北農*42 (8), 13-39 (1975)
- 72) 国井輝男 "上川地方における秋播小麦の冬損に関する研究. 第1報. ほ場における冬損の品種、系統差異" *北農*45(11), 13-24(1978)
- 73) 国井輝男 "同上. 第2報. 越冬前の生育量と冬損とくに越冬茎歩合との関係" *北農*47(6), 1-12(1980)
- 74) 国井輝男 "同上. 第3報. 施肥量とくに基肥窒素の量と冬損について" *北農*47(7), 1-9(1980)
- 75) 国井輝男 "同上. 第4報. 冬損の強弱と秋冬の乾物生産性" *北農*48(5) 1-8(1981)
- 76) 国井輝男 "同上. 第5報. 窒素の分施" *北農*52(8), 1-11 (1985).
- 77) 国井輝男 "同上. 第6報. ほ場における小粒菌核病の人工接種の試み" *育種. 作物学会北海道談話会報*. 26,49(1986)
- 78) 楠隆, 長内俊一 "寒地における秋播小麦の耐寒性検定について" *日作紀*. 23(2), 135(1954)
- 79) 楠隆, 長内俊一, 子安喜代司 "秋播小麦の耐冬性と2、3特性との関係" *北農研抄*. 3, 4-5 (1956)
- 80) 楠隆, 長内俊一, 伊藤平一 "北見地方火山灰地における秋播小麦の冬損を支配する要因についての1実験" *北農研究抄*. 3, 8-9(1956)
- 81) 楠隆, 長内俊一 "秋播小麦雑種初期世代の耐寒性に関する集団選抜の効果" *育種*. 9, 1-6(1959)
- 82) Lamb, C. A. "Physiology. In *Wheat and wheat improvement*" *Amer. Soc. Agron., Inc., Madison, Wis. Press, Agronomy* 13, 181-223 (1967)
- 83) Law, C. N. and G. Jenkins "A genetic study of cold resistance in wheat" *Genet. Res.* 15, 197-208 (1970)
- 84) Levitt, J. "The hardiness of plants" *Agronomy* 6, *Amer. Soc. Agron. Inc., Madison, Wis. Press.* (1956)
- 85) Levitt, J. "Responses of plants to environmental stress." *Academic Press, New York.* 667 pp. (1972)
- 86) Levitt, J. "Responses of plants to environmental stress. Vol. 1. Chilling, freezing, and high temperature stresses" *Academic Press, New York,* 471 p. (1980)
- 87) Li, P. H. and A. Sakai "Plant cold hardiness and freezing stress. Mechanisms and crop implications" *Academic Press, New York,* 416 p. (1981)
- 89) Limin, A. E. and D. B. Fowler "Cold-hardiness response of sequential winter wheat tissue segments to differing temperature regimes" *Crop Sci.* 25, 838-843 (1985)

- 90) Marshall, H. G. "A technique of freezing plant crowns to determine the cold resistance of winter oats" *Crop Sci.* 5, 83-86 (1965)
- 91) Marshall, H. G. "Natural selection for cold resistance in winter oat bulk populations" *Crop Sci.* 13, 581 (1973)
- 92) Marshall, H. G. "Registration of Pennwin oats" *Crop Sci.* 13, 581 (1973)
- 93) Marshall, H. G. and F. L. Kolb "Individual crown selection for resistance to freezing stress in winter oats" *Crop Sci.* 22, 506-510 (1982)
- 94) Marshall, H. G., C. R. Olien and E. H. Everson "Technique for selection of cold hardiness in cereals. In Analysis and improvement of plant cold hardiness" CRC Press, Florida. 139-159 (1981)
- 95) Marshall, H. G. "Breeding for tolerance to heat and cold. In Breeding plants for less favorable environments." John Wiley & Sons, Inc. 47-70 (1982)
- 96) Martin, J. H. "Comparative studies of winter hardiness in wheat" *J. Agric. Res.* 35, 493-535 (1927)
- 97) Matsumoto, N., T. Sato and T. Araki "Biotype differentiation in the *Typhula ishikariensis* complex and their allopatry in Hokkaido" *Ann. Phytopath. Soc. Japan* 48, 275-280 (1982)
- 98) Matsumoto, N., T. Sato "Factors involved in the resistance of timothy and perennial ryegrass to *Sclerotinia borealis* and *Typhula ishikariensis*" *Res. Bull. Hokkaido Agric. Exp.* 136, 23-30 (1983)
- 99) 松本直幸 "雪腐病生物防除についての試論" *北農*52 (11), 1-11 (1985)
- 100) 松尾孝嶺、野村正、"積雪下に於ける小麦の生育に就て" *日作紀.* 12 (4)391-402 (1941)
- 101) 松尾孝嶺、野村正、岩切峻 "農作物の雪害防除に関する試験成績" *農商省農政局*54 p (1944)
- 102) Meryman, H. T., R. J. Williams and M. st. J. Douglas "Freezing injury form solution effect and its prevention by natural or artifitial cryo-protection" *Cryobiology* 14, 287-302 (1977)
- 103) Metcalf, E. L., C. E. Cress, C. R. Olien and E. H. Everson "Relationship between crown moisture content and killing temperature for three wheat and three barley cultivars" *Crop Sci.* 10, 362-365 (1970)
- 104) Muehlbauer, F. I., H. G. Marshall and R. R. Hill, Jr. "Winter hardiness in oat populations derived from reciprocal crosses" *Crop sci.* 10, 646-649 (1970)
- 105) 百足幸一郎 "コムギの雪腐病抵抗性、幼苗検定法の改善" *育雑.* 36 (11), 1797-1800 (1960)
- 106) 中川九一 "麦雪腐病の防除法" *農&園.* 35 (11), 1797-1800 (1960)
- 107) 成田武四 "北海道農作物病害総覧" *北海道植物防疫協会,* 94-107 (1980)
- 108) 能代昌雄、酒井昭 "野草の耐凍性" *日本生態学会誌*24(3), 175-179 (1974)
- 109) 能代昌雄、平島利昭 "牧草の耐凍性に関する研究. I. 北海道根釧地方におけるイネ科牧草の凍害と雪腐大粒菌核病害" *日草誌.* 23 (4), 289-294 (1978)
- 110) 能代昌雄、平島利昭 "根釧地方におけるオーチャードグラス草地の冬枯れ対策法" *日草誌.* 24(4), 277-284 (1979)
- 111) 能代昌雄 "イネ科牧草の凍害と雪腐大粒菌核病発生との関連" *日草誌.* 25 (4), 386-388 (1980)
- 112) 能代昌雄 "牧草の耐凍性に関する研究. II. 2, 3の凍結条件と凍害および増大との関係" *日草誌.* 27, 253-258 (1981)

- 113) Noshiro M, and A. Sakai "Freezing resistance of herbaceous plants" *Low Temp. Sci. Ser. B* 37, 11-18 (1979)
- 114) 農林省農林水産会議事務局 "大型機を中心とした秋播小麦栽培技術体系—北海道東部畑作地帯における—" 地域標準技術体系畑作8, 70 p. (1967)
- 115) 小田桂三郎 "作物体系Ⅱ. 麦の生理, 生態" 養賢堂, 東京(1963)
- 116) 応用統計ハンドブック編集委員会編 "応用統計ハンドブック" 養賢堂東京328-354 (1978)
- 117) Olien, C. R. "Freezing processes in the crown of Hudson barley, *Hordeum vulgale* (L., emd. Lam.) Hudson" *Crop Sci.* 4, 91-95 (1964)
- 118) Olien, C. R. "Influence of cereal polymers and related compounds with freezing" *Cryobiology* 2, 47-54 (1965)
- 119) Olien, C. R. "Freezing stresses and survival" *Ann. Rev. Plant Physiol.* 18, 387-408 (1967)
- 120) Olien, C. R. "Preliminary classification of polysaccharide freezing inhibitors" *Crop Sci.* 7, 156-157 (1967)
- 121) Olien, C. R. "A comparison of desiccation and freezing as stress vectors" *Cryobiology* 8, 244-248 (1971)
- 122) Olien, C. R. "Thermodynamic components of freezing stress" *J. Theor. Biol.* 39, 201-21 (1973)
- 123) Olien, C. R. "Energies of freezing and frost desiccation" *Plant Physiol.* 53, 764-767 (1974)
- 124) Olien, C. R., B. L. Marchetti, and E. V. Chomyn "Ice structure in hardened winter barley" *Mich. Agric. Exp. Stn. Q. Bull.*, 50, 440 (1968)
- 125) Olien, C. R. and S. F. Chao "Liquid water content of cell walls in frozen tissues evaluated by electrophoresis of indicators" *Crop Sci.* 13, 674-676 (1973)
- 127) Olien, C. R. "Barley: patterns of response to freezing stress" *USDA Tech. Bull.* 1558, (1977)
- 128) Olien, C. R. "Analysis of freezing stresses and plant response. In *Plant cold hardiness and freezing stress*" Academic Press, New York. 37-48 (1978)
- 129) Olien, C. R. and M. N. Smith "Protective systems that have evolved in plants. In *Analysis and improvement of plant cold hardiness*" CRC Press, Florida. 61-87 (1981)
- 130) Olien, C. R. "Analysis of midwinter freezing stress. In *Analysis and improvement of plant cold hardiness*" CRC Press, Florida 35-60 (1981)
- 131) 長内俊一 "コムギ生産の安定条件" 日本作物学会シンポジウム記事, 日作紀. 55, 115-143 (1986)
- 132) 尾崎政春 "オーチャードグラス雪腐大粒菌核病の発生生態" 北海道立農試集報. 42, 55-65 (1979) .
- 133) 尾関幸男, 佐々木宏, 天野洋一 "北海道の畑作技術—麦類編—農業技術普及協会, 235p (1981)
- 134) 尾関幸男 "小麦育種の将来と展望" 育種. 作物学会北海道談話会報別刷第2号 第12回シンポジウム資料, (1974) .
- 135) Peltier, G. L. "Control equipment for the study of hardiness in crop plants" *J. Agric. Res.* 43, 177 (1931)
- 136) Poehlman, J. M. and F. A. Coffman "Registration of Dade oats" *Crop Sci.* 9, 296 (1969)
- 137) Pugsley A. T. "The impact of plant physiology on Australian wheat breeding" *Euphytica* 32, 743-748 (1983)
- 138) プルツコーバ, M. T. "冬小麦の品種と耐寒性" 育種と採種4, 23-29 (1970) .
- 146) 齊藤邦八 "小麦の播種期及び幼植物時代の肥料要素欠乏と耐雪性との関係農&園. 14(6), 1527-1533 (1939)

- 139) Quisenberry, K. S. , and J. A. Clark “Breeding hard red winter wheats for winter hardiness and high yield” U. S. Dept. Agric. Tech. Bull. 136,(1929)
- 140) Quisenberry, K. S. “Inheritance of winter hardiness, growth habit and stem rust reaction between Minhardi and H 44 spring wheats” U. S. Dep. Tech. Agric. Bull. 218, (1931)
- 141) Quisenberry, K. S. “Survival of wheat varieties in the Great Plains winterhardiness nursery, 1930–1937” J. Amer. Soc. Agron. 30, 399–405 (1938)
- 142) Quisenberry, K. S. and B. B. Bayles “Growth habit of some winter wheat varieties and its relation to winterhardiness and earliness” J. Amer. Soc. Agron. 31,785–789 (1939)
- 143) Roberts, D. W. A. and M. N. Grant “Changes in cold hardiness accompanying development in winter wheat” Can. J. Plant Sci. 48, 369–376 (1968)
- 144) Roberts, D. W. A. and M. D. MacDonald “Evidence for the multiplicity of alleles at *Vrn 1*, the winter–spring habit locus in common wheat” Can. J. Genet. Cytol. 26, 191–193 (1984)
- 145) Rohde, C. R. and C. F. Pulham “Genetic studies of winter hardiness in barley” Neb. Agric. Exp. Stn. Res. Bull. 193, (1960)
- 147) 酒井昭 “植物の耐凍性と寒冷適応” 学会出版センター,469p. (1982) .
- 148) Salmon, S. C. “Why cereals winterkill” Agron. J. 9, 353–380 (1917)
- 149) Salmon, S. C. “Resistance of varieties of winter wheat and rye to low temperature in relation to winter hardiness and adaptation” Kans. State Univ. Tech. Bull. 35, (1932)
- 150) 佐々木宏 “北海道における小麦育種の現状と問題点” 北海道土壌肥料研究通信第24回シンポジウム特集号,10–25 (1977) .
- 151) Scarth, G. W. “Dehydration injury and resistance” Plant Physiol. 16, 171–179 (1941)
- 152) 進藤武郎,望月太 “ペレニアルグラスの耐寒性の検定. 選抜に関する研究. I. 電気伝導度による耐寒性検定のための測定条件の吟味” 山梨県農試報告. 16,107–174 (1972) .
- 153) Smith, M. N. and C. R. Olien “Pathological factors affecting survival of winter barley following controlled freeze tests” Ecol. Epidemiol. 68,773 (1978)
- 154) Smith, M. N. and C. R. Olien “Toxicity of freeze killed roots as a factor in winter hardiness of barley” Phytopathology 68,773 (1978)
- 155) Steponkus, P. L. “Cold hardiness and freezing injury of agronomic crops” Advances in Agronomy 30, 51–98 (1978)
- 156) Stushnoff, C. , D. B. Fowler and Amta Bruele–Babel “Crop Breeding. V. Breeding and selection for resistance to low temperature” Pergamon Press, 115–136 (1984)
- 157) Sprague, G. F. “Quantitative genetics in crop improvemint. In Plant breeding” K. J. Frey, Ed. Iowa State Univ. Press. 315–354 (1966)
- 158) Sugiyama, N. and T. Simura “Studies on tea varietal differentiation of frost resistance in tea plant. III. With special emphasis on the relation between their frost resistance and chloroplast soluble protein” Japan J. Breed. 17, 37–42 (1967)
- 159) Sugiyama, N. and T. Simura “Studies on the varietal differentiation of frost resistance in the tea plant. IV. The effects of sugar level combined with protein in chloroplasts on frost resistance” Japan J. Breed, 17, 292–296 (1967)
- 160) Suneson, C. A. and G. L. Peltier “Cold resistance ajustments of field hardened winter wheats as determined by artificial freezing” J. Amer. Soc. Agron. 26, 50–58 (1934)

- 161) Suneson, C. A. and G. L. Peltier "Effect of weather variants on field hardening of winter wheat" J. Amer. Soc. Agron. 30, 769-778 (1938)
- 162) Suneson, C. A. and H. Stevens "Studies with bulked hybrid populations of barley" USDA Tech. Bull. No. 1067, 14 pp. (1953)
- 163) Suneson, C. A. "An evolutionary plant breeding method" Agron. J. 48,188-190 (1956)
- 164) Suneson, C. A. and G. A. Wiebe "A 「Paul Bunyan」 plant breeding enterprise with barley" Crop Sci. 2, 347-348 (1962)
- 165) Sutka J. "Genetic studies of frost resistance in wheat" Theor. Appl. Genet. 59, 145-152 (1981)
- 166) Takahashi, R. and S. Yasuda "Genetics of earliness and growth habit in barley" Proc. 2nd. Int. Barley Genet. Symp. 388-408 (1971)
- 167) 滝島英作 "小麦の耐雪性に関する研究. 第1報 積雪下及び高温暗所における生態的变化" 農&園. 15 (4) ,863-870 (1940) .
- 168) 滝島英作 "小麦の耐雪性に関する研究. 第2報 根雪前における発育相の差異と耐雪力(1)" 農&園. 18 (9) ,922-926 (1943) .
- 169) Taylor, J. W. and M. A. McCall "Influence of temperature and other factors on the morphology of the wheat seedling" J. Agric. Res. 52,557-568 (1936) .
- 170) 富山宏平 "麦類雪腐病に関する研究" 北海道農試報告47,1-234 (1955) .
- 171) 富山宏平 "ムギ類雪腐病" 日植病報. 31 (記念号) ,200-206 (1965) .
- 172) 上野賢司, 番場宏治 "秋播小麦における小粒菌核病の人工接種による耐雪性検定について" 育種. 作物学会北海道談話会報. 26,50 (1986) .
- 173) 宇井格生監修 "北海道畑作物の土壤病害" 北海道畑作物の土壤病害刊行会402p. (1983) .
- 174) 宇佐美正一郎 "バーナリゼーション" 植物生理学講座4生活環の制御, 朝倉書店170-186 (1972) .
- 175) Vavilov N. I. "Scientific basis for wheat breeding" Moscow and Leningrad(的場訳) 欧亜通信社(1935)
- 176) Warnes, D. D. , J. W. Schmidt and V. A. Johnson "Correlation of artificial crown-freezing survival with natural survival in winter barley" Barley genetics II, Proc, 2nd. Int. Barley Genet. Symp. , Washington Univ. Press. , 364 (1971)
- 177) Warnes, D. D. and V. A. Johnson "Crown-freezing and natural survival comparisons in winter wheat" Agron. J. 64, 285-288 (1972)
- 178) Webb, R. B. "Crown and root development in wheat seedling" J. Agric. Res. 52, 569-583 (1936)
- 179) Weibel, R. O. and K. S. Quisenberry "Field versus controlled freezing as a measure of cold resistance of winter wheat varieties" J. Amer. Soc. Agron. 33, 336-(1941)
- 180) Wiebe G. A. and D. A. Reid "Comparative winter hardiness of barley varieties" USDA Tech. Bull. No. 1176, (1958)
- 181) Wilner, J. "Relative and absolute electrolytic conductance tests for frost hardiness of apple varieties" Can. J. Plant Sci. 40, 630-637 '1960)
- 182) Worzella, W. W. "Inheritance of cold resistance in winter wheat, with preliminary studies on the technic of artificial freezing tests" J. Agric. Res. 50, 625-638 (1935)
- 183) Worzella, W. W. and G. H. Cutler "Factors affecting cold resistance in winter wheat" J. Amer. Soc. Agron. 33, 221 (1941)
- 184) Worzella, W. W. "Inheritance and interrelationship of components of quality, cold resistance and morphological characters in wheat hybrids" J. Agric. Res. 65, 501-522 (1942)
- 185) Yasuda S. and H. Shimoyama "Analysis of internal factors influencing the heading time of wheat varieties" Ber. Ohara Inst. Landw. Biol. 13, 23-38 (1965)

Studies on Methods of Breeding Wheat for Winter Hardiness

by

Yoichi Amano

Summary

Wheat is the crop most widely grown in the world. Such a broad adaptability is thought to be due mainly to differentialtion of growth habits. Spring wheat can be grown not only in the tropics but also in the Arctic. However in mild climatic areas farmers generally grow winter wheat, wh ch is more productive than spring wheat.

Winter wheat sometimes suffers from serious damage when it is exposed to extremely low temperatures and/or drought conditions, or remains too long under heavy snow cover. So winter hardiness, the ability to survive severe winter climatic conditions, is a main concern in wheat cultivation in such regions.

The causes of winter injury are usually various and complicated. In a particular region any one factor, such as①freezing,②heaving,③smothering,④physiological drought,⑤snow mold may predominate, but in most cases those factors interact with each other. So in breeding for winter hardiness, many factors must be taken into account.

Though Hokkaido is located in the southern part of the subarctic zone, snow persists comparatively long, for 130~170 days. So winter wheat cultivation has long suffered from winter damage. Winter injury in Hokkaido is directly caused by several types of snow mold and in rare cases by freezing, but freezing injury certainly indirectly affects the incidence of snow molds. Tomiyama(1955) clarified that the cause of winter injury was snow molds and developed chemical controls for these diseases. After that, winter damage has decreased remarkably, but in most cases effects of fungicide on these pathogens are still not absolute even in recent cultivation. There are five snow molds, namely *Typhula incarnata*, *T. ishikariensis*, *Fusarium nivale*, *Sclerotinia borealis* and *Pythium* spp. in Hokkaido, and except for *Pythium* spp.,which is localized in badly drained fields, each pathogen occurs over a wide area.

Breeding programs for winter hardiness in Hokkaido have dealt mainly with snow molds and freezing resistance. We have tried to improve the winter hardiness of Hokkaido wheat varieties by introducing snow mold and freezing resistance. In this paper we investigate①development of testing method for winter hardiness and identification of resistant varieties,②genetics of winter hardiness,③characteristics of accumulated substances in resistant varieties,④relationship between growth habits and winter hardiness,⑤ efficiency of selection for winter hardiness. The results obtained are as follows.

1) Varietal differences of 25 winter wheats in winter hardiness

In this experiment several methods for testing *Sclerotinia borealis*, *Typhula ishikariensis*, *Typhula incarnata*, *Fusarium nivale* and freezing injury were devised and using those methods, 25 varieties originating from U.S.S.R., U.S.A., Holland and Hokkaido were tested.

For the method of testing resistance to *S. borealis*, wheat plants were grown on the high ridge

conditions in the fields and sprayed with ascospores suspension of this pathogen. This was more effective than when plants were grown on level ground and in furrow ridge in the fields.

Inoculation of wheat with *Typhula* spp. and *F. nivale* was done in a snow mold chamber using techniques developed by Bruehl et al. (1967). The difference between varieties in resistance to *T. ishikariensis* and *F. nivale*, this inoculation method was applied and for *T. incarnata* a field test was done at Iwamizawa.

Natural exposure(-25°C), artificial crown-freezing(-14°C), and Electric Conductivity(-15°C) methods were applied to test cold hardiness. They were very effective.

As the results of screening for winter hardiness, the following wheat varieties were identified as being useful materials for breeding resistance to the various types of disease and injury. However they are inferior to commercial varieties in other agronomic characteristics.

Iohardi, Hokkai 48, : resistant to *S. borealis*

C. I. 14106, P. I. 172582, P. I. 173438, : resistant to *T. ishikariensis* and *F. nivale*

C. I. 14106, P. I. 173438, Norin 62, : resistant to *T. incarnata*

Lutescens 0329, Valujevskaja, : resistant to freezing injury

None of these materials was found to have *S. borealis* and *Typhula* spp. resistance simultaneously. Correlation of resistance to the above two pathogens in 25 varieties was -0.339 and was not significant. Resistance to *S. borealis* and cold hardiness were closely related. And degrees of resistance to *T. ishikariensis*, *T. incarnata* and *F. nivale* were also highly correlated with each other.

2) Genetics of winter hardiness

In this experiment genetic mechanisms for resistance to *S. borealis*, *T. ishikariensis*, *T. incarnata* and freezing injury were analyzed. Diallel crosses were carried out among 7 varieties which had typical levels of resistance to those factors, and parents and F₂ population were examined under each testing method respectively, as mentioned in the previous chapter. A randomised block design with 2 to 10 replications was used and resistance was expressed by the degree of injury.

According to analysis of the variance of means and the variance, differences between parents and F₂ populations were significant in every factor. The correlations of mean and the variance between F₂ populations and midparents were also highly significant. So the resistance of progeny was very likely to be influenced by that of the parent.

Variance analyses of W_r-V_r among replications and tests of regression of W_r to V_r confirmed the validity of the assumption of independent action of non-allelic genes. The hypothesis of diallel analysis was satisfied, so that (V_r, W_r) graphical analysis was done and the genetic variance components and parameters for each factor were estimated according to the method of Hayman and Jinks.

In every factor the additive effects of genes were significantly large, but the dominance effects differed according to each. Resistance to *S. borealis* had partial dominance and resistance to *T. ishikariensis* and freezing injury showed overdominant. The position of array points along the line of regression showed that resistant parents of *S. borealis* and *T. ishikariensis* had more recessive alleles. On the other hand, a clear tendency of frequencies of effective alleles did not exist in cold

hardiness. In the case of *T. ishikariensis*, approximately equal numbers of dominant and recessive alleles were present, but dominant and recessive alleles were not distributed equally in the cases of *S. borealis* and cold hardiness.

The heritabilities in a narrow sense of *S. borealis*, *T. ishikariensis* and freezing injury were generally high, namely 90, 79, and 82 %, respectively. The number of alleles concerned with each factor was estimated to be 2 to 3, based on the formula, (maximum parent value–minimum parent value)²/4D.

Genetic correlation of the mean of F₂ populations between *S. borealis* and *T. ishikariensis* was highly negative. This suggested that breeding for resistance to both *S. borealis* and *T. ishikariensis* would be laborious work. F₂ populations of the cross between varieties bred in Hokkaido contained more non-damaged plants than the other crosses, so it was surmised that Hokkaido varieties, Norin 8, Muka, Horoshiri and Kitakei 628 had the more stable and satisfactory genetic backgrounds for winter hardiness. Resistant genes are also available in other varieties, such as P. I. 173438 and Moscow 1, so it seems possible to be able to improve the genetic background of winter hardiness in Hokkaido varieties by introducing specific resistant genes for other factors.

3) Relationship between accumulated substances and winter hardiness

In this experiment we dealt with regional and varietal differences of growth rates and accumulations or depletions of substances during winter, and relationships between reserve substances and resistances to freezing, *S. borealis*, *Typhula* spp., and *F. nivale*. The 25 varieties mentioned in chapter I were examined at Kunneppu and Iwamizawa in 1978-'79 and 1979-'80 seasons. Twenty characteristics measured four times before and after winter were used for analysis. The degree of resistance to freezing and snow molds obtained in previous experiments were taken into account.

Dry weight, percentage of dry matter, total carbohydrate, sugar, starch, and proteinous nitrogen in the shoot of the plant increased rapidly from autumn to early winter, and reached the maximum amount immediately after the beginning of permanent snow. Soluble nitrogen, phosphoric acid and silic acid, decreased with the decrease of temperature. Differences between locations, seasons, varieties, and wintering types shown in chapter I were significant in 16 characteristics measured before winter. Wheat plants in Kunneppu showed small growth rate, high accumulations of carbohydrate and sugar in the fall, and small depletions of them during winter than wheat plants in Iwamizawa. Varieties resistant to *Typhula* spp. and *F. nivale* entered winter with greater carbohydrate reserves and possessed much more of them after snow melts than susceptible varieties.

Correlations of reserve substances levels before and after snow cover were highly significant except for reducing sugar and starch. Correlation coefficients of total carbohydrate between years was 0.751*** (before winter) and 0.767*** (after snow melt) at Kunneppu. Highly significant correlations were also found between following resistances and characters. Freezing and osmotic value; *S. borealis* and phosphoric acid, lipid; *Typhula* spp., *F. nivale* and proteinous nitrogen, total carbohydrate.

Varieties were classified into four groups based on the principal component analysis using 22

characteristics of plant size and reserve substances. These groups coincided very well with the above mentioned wintering groups, namely, A(less hardy), B(cold resistant), C(intermediate), D(snow endurable). Resistant varieties to *Typhula* spp. and *F. nivale*, 「C. I. 14106」, 「P. I. 173438」 selected by Bruehl et al. accumulated more carbohydrate more rapidly in the fall than other varieties did. Nevertheless, their content of proteinous nitrogen, phosphoric acid, lipid, and their osmotic value were very low. These facts may be a reason why the above three varieties show susceptibility to freezing and *S. borealis*. These characteristics might be ascribed to genetic specificity.

4) Growth habits and winter hardiness

The 26 winter wheat varieties mentioned in chapter I and 「Chihoku」 were divided into four groups according to the degree of vernalization requirement(DVR). Among them a new group was recognized, which requires longer periods of vernalization compared to the VII group designated as an extreme winter habit growth type and/or DVR. In this group cold hardy varieties (Moscow 1, Valujevskaja, and Lutescens 0329) and snow mold resistant varieties (C. I. 14106, P. I. 172582. and P. I. 173438) were included. In order to identify DVR of a high level as with these varieties, treatment with short-day photoperiods and low temperatures(0~2°C) were more effective than that of continuous illumination under comparatively high temperatures(8°C) for green plants.

Wheat lines with various high groups of DVR were developed from two crosses of very high DVR varieties (P. I. 173438 and Valujevskaja) with a variety of DVRVII(Chihoku). In each level of DVR, wheat lines having resistance to cold injury or *Typhula ishikariensis* were found. It was suggested that high DVR traits are under the control of several genes and their actions are independent of cold hardiness and snow mold resistance, because there are not any correlations between DVR and cold hardiness or resistance to *S. borealis* and *Typhula* spp. in the 26 varieties nor in the many lines obtained from the crosses mentioned above.

5) Efficiency of selection for winter hardiness

To improve the winter hardiness of Hokkaido wheat, we intended to introduce resistant genes for two traits. One is resistance to *Typhula* spp, and *F. nivale* represented by 「C. I. 14106」, 「P. I. 173438」, and the other is freezing resistance represented by 「Valujevskaja」, 「Moscow 1」. For that object we operated series of selection experiments.

(1) Series 1 : (Cross 1051 : P. I. 173438/Horoshiri//Kitami42)

We performed a selection in this cross using pedigree method from F₂ to F₄ at Kitami where *S. borealis* usually occur. We also referred to the results of snow mold tests conducted at Shibetsu, where incidence of snow molds, especially *Typhula* spp. is high. In F₅, progeny lines were exposed to natural selection at Sapporo where *Typhula* spp. usually occurs. In F₆, we retreated 91 selected lines with *T. ishikariensis* inoculation in the field at Kitami. The inoculation of sclerotia of *T. ishikariensis* was effective in the field at Kitami. Finally 4 prospective lines which are resistant to *Typhula* spp. and have fairly good agronomic characteristics were identified.

(2) Series 2 : (Cross 1050 : Horoshiri/P. I. 173438//Valujevskaja/Horoshiri)

Hybrid populations were grown as a bulk during F₂ and F₃ generation at Shibetsu for natural

selection to winter hardiness. 20 and 15 percent of plants were killed by *Typhula* spp. in each generation. In F₄, 3,000 plants were planted at Kitami, but there was no winter kill. So 980 plants were sampled randomly. In F₅, 980 lines were treated with *T. ishkariensis* inoculation and freezing tests, respectively. 76 lines (7.8%) out of 980 were recognized as lines resistant to *T. ishkariensis* and 91 lines (9.3%) were as hardy as 'Valujevskaja' in freezing resistance. Six lines (0.6%) were identified as the materials resistant to both traits, but these selected lines were inferior to 'Horoshirikomugi' in other agronomic traits.

(3) Series 3: (Cross 1: Moscow 1/Horoshiri, Cross 2: Muka/Horoshiri, Cross 3: C. I. 14106/Horoshiri)

Hybrid populations were grown as a bulk in F₂, F₃ and F₄ generation at Kitami. During these generations winter kill was very rare and in F₄, 184(Cross 1), 161(2) and 177(3) plants were sampled randomly in each cross. From F₅ to F₇ generations those lines were treated with freezing, *Typhula* spp. and *S. borealis* tests. In every trait differences in resistance among crosses were significant and the average degrees of damage in each trait were as follows,

Freezing(E. C. value) : Cross 3(497) > Cross 2(438) > Cross 1(393)

S. borealis(%) : Cross 3(62) > Cross 2(48) > Cross 1(43)

Typhula spp.(%) : Cross 1(68) > Cross 2(57) > Cross 3(31)

Five prospective lines, three being resistant to *S. borealis* from cross 1 and 2, and two being resistant to *Typhula* spp. from cross 3, were identified. These selected lines were inferior to 'Horoshirikomugi' in other agronomic characteristics.

From the materials of these three series, following 12 lines were identified as genetic resources for winter hardiness.

Resistant to *T. ishkariensis* : WH 3, 5, 6, 7, 8, 9, 10.

Hardy to freezing : WH 1, 2, 3, 4, 6, 11, 12.

Resistant to *S. borealis* : WH 11, 12.

The degree of resistance to *S. borealis* of above mentioned two lines are not seemed to be satisfactory under actual practices, so further investigations are necessary for getting good genetic resources for that trait.