

北海道立総合研究機構 農業試験場報告

第 147 号

Female sex pheromone in the rice leaf bug, *Trigonotylus caelestialium* (Kirkaldy) (Heteroptera: Miridae):
its identification and its use for control

アカヒゲホソミドリカスミカメ（カメムシ目：カスミカメムシ科）の
性フェロモンの解明と防除への利用に関する研究

平成31年3月

北海道立総合研究機構 中央農業試験場

**Female sex pheromone in the rice leaf bug, *Trigonotylus caelestialium* (Kirkaldy)
(Heteroptera: Miridae): its identification and its use for control ***

アカヒゲホソミドリカスミカメ（カメムシ目：カスミカムシ科）の性フェロモンの解明
と防除への利用に関する研究

博士(農学) 柿崎昌志**

Contents

Chapter I GENERAL INTRODUCTION	1
Chapter II Attraction of males to females in the rice leaf bug, <i>Trigonotylus caelestialium</i>	3
2.1 INTRODUCTION	3
2.2 MATERIALS AND METHODS	3
2.2.1 Observation of Mating Behavior	3
2.2.2 Insect-Baited Traps	3
2.3 RESULTS AND DISCUSSION	4
2.3.1 Mating Behavior	4
2.3.2 Trapping Tests in the Spring Wheat Field	5
Chapter III Identification of the female sex pheromone of the rice leaf bug, <i>Trigonotylus caelestialium</i>	7
3.1 INTRODUCTION	7
3.2 METHODS AND MATERIALS	7
3.2.1 Insects and Pheromone Extraction	7
3.2.2 Column Chromatography	7
3.2.3 Bioassay by Coupled Gas Chromatography-Electroantennographic Detector (GC-EAD)	7
3.2.4 Chemical Analysis	7
3.2.5 Chemical Synthesis	8
3.2.6 Baits	8
3.2.7 Field Experiments	8
3.3 RESULTS	9
3.3.1 EAD Active Peaks in the Extracts	9
3.3.2 Identification of the EAD Active Compounds	9
3.3.3 Ratio of the Compounds in Extracts	9
3.3.4 Attraction Activity of EAD Active Compounds in the Field	11
3.3.5 Optimum Ratio and Doses of the Three Compounds for Male Attraction	13
3.3.6 Comparison of Numbers of Bugs Captured by the Synthetic Sex Pheromone and Two Virgin Females	13
3.4 DISCUSSION	15

Chapter IV Mating disruption using the sex pheromone of the rice leaf bug, <i>Trigonotylus caelestialium</i>16
4.1 INTRODUCTION16
4.2 MATERIALS AND METHODS16
4.2.1 Pheromone compounds16
4.2.2 Bait and trap16
4.2.2.1 Pheromone lure16
4.2.2.2 Virgin females16
4.2.2.3 Traps16
4.2.3 Dispensers for mating disruption16
4.2.4 Laboratory tests in a rearing cage17
4.2.5 Field experiments17
4.2.5.1 Small-scale field tests17
4.2.5.2 Large-scale field tests17
4.2.6 Statistics18
4.3 RESULTS18
4.3.1 Release rate of sex pheromone from dispensers for mating disruption18
4.3.2 Laboratory experiment20
4.3.2.1 Effect of the 3-component sex pheromone on copulation20
4.3.2.2 Effect of the 3-component sex pheromone on reproduction20
4.3.3 Disruption of male trapping by sex pheromone components in field experiments21
4.3.4 Population suppression by treatment with the 3-component sex pheromone in field experiments	...23
4.4 DISCUSSION26
Chapter V GENERAL DISCUSSION28
5.1 The female sex pheromone28
5.2 Monitoring and mating disruption using the sex pheromone29
Summary31
和 文 要 旨32
ACKNOWLEDGMENTS34
REFERENCES35

*東京農工大学連合大学院審査学位論文

**北海道立総合研究機構中央農業試験場 (〒069-1395 北海道夕張郡長沼町東6線北15号)

ISSN 2186-1064

北海道立総合研究機構 農業試験場報告 第 147 号

Female sex pheromone in the rice leaf bug, *Trigonotylus caelestialium*
(Kirkaldy) (Heteroptera: Miridae):
its identification and its use for control

(アカヒゲホソミドリカスミカメ（カメムシ目：カスミカメムシ科）の
性フェロモンの解明と防除への利用に関する研究）

著 者 柿 崎 昌 志

平成 31 年 3 月 15 日

発行者 北海道立総合研究機構 中央農業試験場

069-1395 北海道夕張郡長沼町東 6 線北 15 号

印刷所 北海道印刷企画株式会社

REPORT
OF
HOKKAIDO PREFECTURAL AGRICULTURAL
EXPERIMENT STATIONS
No.147

Female sex pheromone in the rice leaf bug, *Trigonotylus*
caelestialium (Kirkaldy) (Heteroptera: Miridae):
its identification and its use for control

by
Masashi Kakizaki

Published by
Hokkaido Research Organization
Central Agricultural Experiment Station
Naganuma, Hokkaido, 069-13, Japan

2019